US ERA ARCHIVE DOCUMENT

#### DATA EVALUATION RECORD

- Chemical: Sulfosate (Shaughnessy No. 128501) 1.
- SC-0224 4-LC (liquid Formulated product: 2. Test Material: concentrate)

40.8% ai (as per proposed label of study sponsor)

Fish acute LC50 Study/Action Type: 3:

Rainbow trout (Salmo gairdneri)

The acute toxicity of SC-0224 4-LC to the Rainbow Study ID: trout (Salmo gairdneri)

> Biospherics, Inc. Project No. 82-E-1119R Report date: November 1982

Study sponsor: Stauffer Chemical Company Study location: EPA Accession No. 250545

James D. Felkel Reviewed by: 5.

Wildlife Biologist

EEB/HED

Signature: James J. Flish

Date: 1-20-87

Signature: Henry Thouse

Date: 1/2487

Approved by: Henry Craven

Supervisory Biologist

EEB/HED

7. Conclusions:

This study is scientifically sound. With a 96-hr  $LC_{50}$ of 5.7 mg/L, this formulation is considered "moderately toxic" to the rainbow trout. A data requirement for this study may be fulfilled with submission of photoperiod information and confirmation of exact formulation composition (i.e., whether exactly as shown in June 3, 1983 Confidential Statement of Formula).

Recommendations: 8.

> The study may be upgraded from supplemental to core for this formulation with the information cited in #7 above.

- Background: N/A. 9.
- Discussion of Individual Test: N/A. 10.

#### 11. Materials and Methods:

- a. Test animals Rainbow trout (Salmo gairdneri) from Trout Lodge, Washington State; mean length 45.2 mm (40.2 to 47.8 mm); mean weight 0.61 g (0.43 to 0.81 g); approximately 2 months old.
- b. Dosage form In static bioassay, no solvent.
- .c. Study Design
  - (1) Range finding: polystyene containers
     4 L test solution
     4 fish/vessel
     0.1, 1.0, 10, 100, 1000 mg/L
     96 hr

  - (3) Statistics Binomial probability method (Stephan 1979).

source - 400' well

#### 12. Reported Results:

```
DO - 5.6 to 9.3 mg/L
pH - 7.0 to 7.5
Mean temp. - 13.0 °C
                              > 8 mg/L
               24-hr LC50
               48-hr LC50
                                  7 mg/L (4-infinity)
                72-hr LC<sub>50</sub>
                                  6 \text{ mg/L} (4-8 \text{ mg/L})
                96-hr LC<sub>50</sub>
                                  6 \text{ mg/L} (4-8 \text{ mg/L})
                24-hr NOEL
                                  4 mg/L
                48-hr NOEL
                                  2 mg/L
                72-hr NOEL
                                  2 \text{ mg/L}
                                  2 mg/L
               96-hr NOEL
```

## 13. Study Author's Conclusions/QA Measures:

See above reported results. Test "... inspected by the Quality Assurance Unit according to Biospherics SOP 2.3.4 'QA Study Inspections'."

## 14. Reviewer's Discussion and Interpretation of the Study:

a. <u>Test Procedure</u>: Polystyrene containers used in the range-finding test would not be acceptable in the definitive study.

Reported procedures for the definitive test were generally consistent with current guidelines (1982). However, photoperiod was not reported and water temperatures were measured daily rather than continuously as recommended by the Committee on Methods for Toxicity Tests with Aquatic Organisms (1975). Also, the 0.6-1-2-4-8-mg/L test series is not geometric, nor are all levels  $\geq$  60% of the next level. Dilution water was harder than recommended (115 mg/L as CaCO<sub>3</sub>).

- b. Statistical Analysis: A check of the statistical analysis was conducted and is attached.
- Discussion/Results: An approximate LC50 is determined to be 5.7 mg/L, very close to that reported in the study. The dose-response curve is very steep, with no mortality at 4 mg/L, but 100% mortality at 8 mg/L. The formulated product appears to be approximately 316X as toxic as technical material to the rainbow trout (technical LC50 with rainbow trout reported in August 9, 1983 EEB review: 1800 mg/L).

### d. Adequacy of the Study:

- (1) Classification: Supplemental for this formulation.
- (2) Rationale: While the study is scientifically sound, the photoperiod should be reported and the registrant must confirm whether the composition of formulation is exactly as described in the June 3, 1983 Confidential Statement of Formula.
- (3) Reparability: Study may be upgraded to core for this formulation with submission of information cited above.

# 15. Completion of One-Liner for Study:

One-liner generated automatically by current computer program.

16. CBI Appendix: N/A.

#### J. FELKEL SC-0224 4-LC RAINBOW TROUT 12-15-86

****	******	****		
CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB. (PERCENT)
8	10	10	100	9.765625E-02
4	10	0	0	9.765625E-02
2	10	0	0	9.765625E-02
1	10	Ó	0	9.765625E-02
.6.	10	1	10	1.074219

THE BINOMIAL TEST SHOWS THAT 4 AND 8 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS, BECAUSE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 5.656854.

WHEN THERE ARE LESS THAN TWO CONCENTRATIONS AT WHICH THE PERCENT DEAD IS BETWEEN 0 AND 100. NEITHER THE MOVING AVERAGE NOR THE PROBIT METHOD CAN GIVE ANY STATISTICALLY SOUND RESULTS.

\*\*\*\*\*\*\*\*\*\*\*\*\*\*